

Anti - PRION

Rabbit clonal antibody

CAT#

DB 081-0.05 (50 µl)
DB 081-0.1 (100 µl)

PRODUCT INFORMATION


Clone number:	C16-S	Buffer:	20 mM Tris-HCl, pH 8.0
Uniprot:	Human: P04156; Ovine: P23907; Bovine: P10279; Mouse: P04925	Stabilizer:	10 mg/ml BSA
Product description:	Rabbit anti-prion clonal IgGs	Preservative:	0.05% Sodium Azide
Basic information:	Major clone of rabbit immunoglobulin corresponding to immunogenic peptide; antibody recognizes PrP ^C , with the potential fo PrP ^{SC} detection	Storage:	10 µl aliquots at -20°C
Immunogen:	Peptide derived from C-terminal sequence of third alpha-helical domain of human prion protein. Antibody recognizes the epitope located between Cys214 - Gly229.	Handling:	Avoid repeated freezing and thawing
Species Reactivity:	Human, ovine, bovine, mouse - tested	Expiration:	24 months from the shipping date
		Applications:	Western blot, Immunoprecipitation, ELISA
		Dilution range:	Western blotting – 1:2,000 ELISA – 1:20,000 – 1:50,000 Immunoprecipitation – dilutions to be tested by user.

WESTERN BLOT (WB) PROTOCOL - INSTRUCTION MANUAL

Western immunoblotting solutions:

- Wash buffer: 1x Tris Buffered Saline (TBS); 0.1% Triton X-100
- Blocking buffer: 1xTBS; 0.2% Tween 20; 8% nonfat dry milk

For western blots, incubate the membrane with antibody diluted in blocking buffer for 2 hours at room temperature.



Anti-prion (DB 081)

- Western blot analysis of prion protein in mouse brain extract (A), HEK293 cells producing recombinant human prion protein (B), and untransfected HEK 293 cells (C). 200 µg of total protein loaded per well.
- Western blot analysis of prion protein was performed by Drs. Valeriy Ostapchenko and Marco Prado, Robarts Research Institute, the University of Western Ontario, London, Ontario, Canada. Lanes A-C represent extracts (30 µg of total protein per lane) of PrP-KO CF-10 cells (A), mouse hippocampus (B) and HEK293 cells, transfected with mouse prion protein bearing 3F4 epitope (C); lane D corresponds to 30 ng recombinant MoPrP.

PRECAUTIONS

- Intended for professional In Vitro Diagnostic use in laboratories.
- Do not use after expiration date stamped on vial label.
- Avoid contamination of the reagent.
- Any discrepancies in the recommended procedures stated in the working protocol may affect the final results.
- The reagent contains sodium azide (NaN₃) which is highly toxic in higher concentrations. The concentration in the reagent (0.05%) is not considered as hazardous.
- Disposal of waste material must be conducted in accordance with local regulations.
- Wear appropriate Personal Protective Equipment to avoid contact with eyes and skin.