

Anti - p38- γ (MAPK12) *Rabbit clonal antibody*

CAT#

DB 163-0.05 (50 μ l)
 DB 163-0.1 (100 μ l)

PRODUCT INFORMATION

Clone number: R13-L
Uniprot: Human: P53778, Mouse: O08911, Rat: Q63538
Product description: Rabbit anti-p38- γ clonal IgGs
Basic information: Major clone of rabbit immunoglobulin corresponding to immunogenic peptide
Immunogen: Peptide derived from the C-terminal sequence of human p38- γ . Antibody recognizes the epitope located between Arg355 - Pro366.
Species Reactivity: Human, mouse, rat - tested


Buffer: 20 mM Tris-HCl, pH 8.0
Stabilizer: 10 mg/ml BSA
Preservative: 0.05% Sodium Azide
Storage: 10 μ l aliquots at -20°C
Handling: Avoid repeated freezing and thawing
Expiration: 24 months from the shipping date
Applications: Western blot, Immunoprecipitation, ELISA, Immunocytochemistry (ICC)
Dilution range: Western blotting – 1:1,000
 ELISA – 1:10,000-1:20,000
 Immunoprecipitation – to be tested by user

WESTERN BLOT (WB) PROTOCOL - INSTRUCTION MANUAL

Western immunoblotting solutions:

- Wash buffer: 1x Tris Buffered Saline (TBS); 0.1% Triton X-100
 - Blocking buffer: 1xTBS; 0.1% Triton X-100; 5% nonfat dry milk

For western blots, incubate the membrane with antibody diluted in blocking buffer for 2 hours at room temperature.




Anti - p38- γ (DB 163)

Western blot analysis of p38- γ . Lane A – human recombinant p38- α (260-360a.a., N-terminal GST tag; Novus Biologicals, cat. #: H00001432-Q01), lane B – human recombinant p38- β (with C-terminal DDK tag, Novus Biologicals, cat. #: NBL1-12870), lane C - human recombinant PKC- γ (1-368a.a., N-terminal GST tag, Novus Biologicals, cat. #: H00006300-P01), lane D – human recombinant p38- δ (1-366a.a., N-terminal GST tag, Novus Biologicals, cat. #: H00005603-P01). 500 ng of recombinant protein was loaded in each well.

IMMUNOCYTOCHEMISTRY (ICC) PROTOCOL - INSTRUCTION MANUAL

- Coat coverslips with 1% gelatin-coating solution for 2 hours at room temperature (RT); rinse with distilled water, and let to dry overnight. Before plating the cells, wash the coated coverslips briefly with PBS.
- Fix the cells with 4% paraformaldehyde solution (in PBS, pH 7.2), for 15 min at RT.
- Wash 2 x 3 min with PBS.
- Permeabilize the cells with 0.1% Triton X-100 solution (in PBS, pH 7.2) for 5 min on ice.
- Wash 2 x 3 min with PBS.
- Incubate the cells in blocking buffer (0.3M glycine in PBS, 2% BSA) for 30 min at RT.
- Incubate the cells with primary antibody: anti-p38- γ clonal antibody at the dilution of **1:200 - 1:500** in antibody dilution buffer (PBS, 1% BSA) for 1 hour at RT in humid chamber.
- Wash 2 x 3 min with PBS.
- Apply the secondary antibody (*in this case, the goat anti-rabbit IgG-FITC from Jackson ImmunoResearch, cat. # 111-095-003, was used* at 1:300 in antibody dilution buffer, and cells were incubated for 1 hour at RT in dark).
- Wash 3 x 3 min with PBS.
- Rinse once with distilled water.
- Mount the slide for observation, with a drop of anti-fade mounting medium.



Representative picture of p38- γ expression in HEK293 cells, visualized with clonal rabbit anti-p38- γ monospecific antibody. Primary antibody dilution - 1:300.

PRECAUTIONS

- Intended for professional In Vitro Diagnostic use in laboratories.
- Do not use after expiration date stamped on vial label.
- Avoid contamination of the reagent.
- Any discrepancies in the recommended procedures stated in the working protocol may affect the final results.
- The reagent contains sodium azide (NaN₃) which is highly toxic in higher concentrations. The concentration in the reagent (0.05%) is not considered as hazardous.
- Disposal of waste material must be conducted in accordance with local regulations.
- Wear appropriate Personal Protective Equipment to avoid contact with eyes and skin.