

Anti - Akt2 (pSer478)

Rabbit clonal antibody

CAT#

DB 183-0.05 (50 µl)
DB 183-0.1 (100 µl)


PRODUCT INFORMATION

Clone number: Q10-E
Uniprot: Human: P31751; Mouse: Q60823; Rat: P47197
Product description: Rabbit anti-AKT2 (pSe478) clonal IgGs
Basic information: Major clone of rabbit immunoglobulin corresponding to immunogenic peptide
Immunogen: Peptide surrounding pSer-478 at the C-terminal sequence of human Akt2
Species Reactivity: Human, mouse, rat - tested

Buffer: 20 mM Tris-HCl, pH 8.0
Stabilizer: 10 mg/ml BSA
Preservative: 0.05% Sodium Azide
Storage: 10 µl aliquots at -20°C
Handling: Avoid repeated freezing and thawing
Expiration: 24 months from the shipping date
Applications: Immunocytochemistry (ICC)
Dilution range: 1:500 – 1:1000

IMMUNOCYTOCHEMISTRY (ICC) PROTOCOL - INSTRUCTION MANUAL

1. Coat coverslips with 1% gelatin-coating solution for 2 hours at room temperature (RT); rinse with distilled water, and let to dry overnight. Before plating the cells, wash the coated coverslips briefly with PBS.
2. Fix the cells with 4% paraformaldehyde solution (in PBS, pH 7.2), for 15 min at RT.
3. Wash 2 x 3 min with PBS.
4. Permeabilize the cells with 0.1% Triton X-100 solution (in PBS, pH 7.2) for 5 min on ice.
5. Wash 2 x 3 min with PBS.
6. Incubate the cells in blocking buffer (0.3M glycine in PBS, 2% BSA) for 30 min at RT.
7. Incubate the cells with primary antibody: anti-Akt2, pSer478 clonal antibody at the dilution of **1:500 - 1:2000** in antibody dilution buffer (PBS, 1% BSA) for 1 hour at RT in humid chamber.
8. Wash 2 x 3 min with PBS.
9. Apply the secondary antibody (*in this case, the goat anti-rabbit IgG-TRITC from Jackson Immunoresearch, cat. # 111-025-003, was used at 1:200 in antibody dilution buffer, and cells were incubated for 1 hour at RT in dark.*)
10. Wash 3 x 3 min with PBS.
11. Rinse once with distilled water.
12. Mount the slide for observation, with a drop of anti-fade mounting medium.



Representative pictures of pS478-Akt2 expression in Akt2 transfected and PMA stimulated HEK293 cells (A); and in non-transfected, untreated HEK293 cells (B), visualized with anti-pS478-Akt2 rabbit monospecific clonal antibody. Primary antibody dilution - 1:1000.

PRECAUTIONS

1. Intended for professional In Vitro Diagnostic use in laboratories.
2. Do not use after expiration date stamped on vial label.
3. Avoid contamination of the reagent.
4. Any discrepancies in the recommended procedures stated in the working protocol may affect the final results.
5. The reagent contains sodium azide (NaN₃) which is highly toxic in higher concentrations. The concentration in the reagent (0.05%) is not considered as hazardous.
6. Disposal of waste material must be conducted in accordance with local regulations.
7. Wear appropriate Personal Protective Equipment to avoid contact with eyes and skin.