

Anti - p38-β (MAPK11)

Rabbit clonal antibody

CAT#

DB 162-0.05 (50 µl)
DB 162-0.1 (100 µl)

PRODUCT INFORMATION


Clone number:	E13-Q	Buffer:	20 mM Tris-HCl, pH 8.0
Uniprot:	Q15759	Stabilizer:	10 mg/ml BSA
Product description:	Rabbit anti-p38-β clonal IgGs	Preservative:	0.05% Sodium Azide
Basic information:	Major clone of rabbit immunoglobulin corresponding to immunogenic peptide	Storage:	10 µl aliquots at -20°C
Immunogen:	Peptide derived from the C-terminus of human p38-β sequence. Antibody recognizes the epitope located between Glu352 - Glu363.	Handling:	Avoid repeated freezing and thawing
Species Reactivity:	Human specific - tested	Expiration:	24 months from the shipping date
		Applications:	Western blot, Immunoprecipitation, ELISA, Immunocytochemistry (ICC)
		Dilution range:	Western blotting – 1:1,000 ELISA – 1:10,000-1:20,000 Immunoprecipitation – to be tested by user

WESTERN BLOT (WB) PROTOCOL - INSTRUCTION MANUAL

Western immunoblotting solutions:

- Wash buffer: 1x Tris Buffered Saline (TBS); 0.1% Triton X-100
- Blocking buffer: 1xTBS; 0.1% Triton X-100; 5% nonfat dry milk

For western blots, incubate the membrane with antibody diluted in blocking buffer for 2 hours at room temperature.




Anti - p38-β (DB 162)

Western blot analysis of p38-β. Lane A – human recombinant p38-α (260-360a.a., N-terminal GST tag; Novus Biologicals, cat. #: H00001432-Q01), lane B – human recombinant p38-β (with C-terminal DDK tag, Novus Biologicals, cat. #: NBL1-12870), lane C - human recombinant PKC-γ (1-368a.a., N-terminal GST tag, Novus Biologicals, cat. #: H00006300-P01), lane D – human recombinant p38-δ (1-366a.a., N-terminal GST tag, Novus Biologicals, cat. #: H00005603-P01). 500 ng of recombinant protein was loaded in each well.

IMMUNOCYTOCHEMISTRY (ICC) PROTOCOL - INSTRUCTION MANUAL

1. Coat coverslips with 1% gelatin-coating solution for 2 hours at room temperature (RT); rinse with distilled water, and let to dry overnight. Before plating the cells, wash the coated coverslips briefly with PBS.
2. Fix the cells with 4% paraformaldehyde solution (in PBS, pH 7.2), for 15 min at RT.
3. Wash 2 x 3 min with PBS.
4. Permeabilize the cells with 0.1% Triton X-100 solution (in PBS, pH 7.2) for 5 min on ice.
5. Wash 2 x 3 min with PBS.
6. Incubate the cells in blocking buffer (0.3M glycine in PBS, 2% BSA) for 30 min at RT.
7. Incubate the cells with primary antibody: anti-p38-β clonal antibody at the dilution of **1:400 - 1:1000** in antibody dilution buffer (PBS, 1% BSA) for 1 hour at RT in humid chamber.
8. Wash 2 x 3 min with PBS.
9. Apply the secondary antibody (*in this case, the goat anti-rabbit IgG-FITC from Jackson ImmunoResearch, cat. # 111-095-003, was used at 1:300 in antibody dilution buffer, and cells were incubated for 1 hour at RT in dark*).
10. Wash 3 x 3 min with PBS.
11. Rinse once with distilled water.
12. Mount the slide for observation, with a drop of anti-fade mounting medium.



Representative picture of p38-β expression in HEK293 cells, visualized with clonal rabbit anti-p38-β monospecific antibody. Primary antibody dilution - 1:500.

PRECAUTIONS

1. Intended for professional In Vitro Diagnostic use in laboratories.
2. Do not use after expiration date stamped on vial label.
3. Avoid contamination of the reagent.
4. Any discrepancies in the recommended procedures stated in the working protocol may affect the final results.
5. The reagent contains sodium azide (NaN₃) which is highly toxic in higher concentrations. The concentration in the reagent (0.05%) is not considered as hazardous.
6. Disposal of waste material must be conducted in accordance with local regulations.
7. Wear appropriate Personal Protective Equipment to avoid contact with eyes and skin.